

REPLACED BY
ART 34 AMDT

Claims

✓
1. The method of synthesizing phenstatin comprising the steps of:
oxidizing 3-(tert-butyl dimethylsilyl)oxy-4-methoxybenzaldehyde with
potassium permanganate to form the corresponding carboxylic acid;

5
Sub. B1
converting said carboxylic acid to the corresponding acid chloride;

10
treating said acid chloride with the lithium derivative obtained from 3,
4, 5-trimethoxybenzene and t-butyllithium to form a protected product;

and deprotecting said protected product to form phenstatin.

15
2. The method of synthesizing phenstatin prodrug comprising the steps of:


phosphorylating phenstatin with dibenzylphosphite in the presence of
bromodichloromethane to form a phosphate ester;

20
cleaving the benzyl groups from said phosphate ester by means of
catalytic hydrogenolysis; and

reacting the cleaved phosphate ester with sodium methoxide to produce
the phenstatin sodium phosphate prodrug.


25
3. The method of inhibiting cancer cell growth and tubulin polymerization
in an environment inflicted therewith comprising: introducing into said
environment a pharmaceutically acceptable carrier and a small but effective
amount of phenstatin prodrug.

4. Phenstatin prodrugs and derivatives thereof having the structure:





wherein when $R=H$ and $R_1 = OCH_3$, R_2 is OPO_3Na_2 , $OCOCH_3$, H , or OCH_3 and when $R=R_2$, R_2 is OCH_3 , CH_3 , CL or F and R_1 is H and when $R_1=R_2$, R_2 is OCH_3 or OCH_2O and R is H .

5. The method of inhibiting human cancer cell growth in a host inflicted therewith comprising administering to said host in pharmaceutically acceptable carrier a small but effective amount of a compound selected from the group consisting of phenstatin, phenstatin prodrug and the derivatives thereof having the structure.




wherein when $R=H$ and $R_1 = OCH_3$, R_2 is OPO_3Na_2 , $OCOCH_3$, H , or OCH_3 and when $R=R_2$, R_2 is OCH_3 , CH_3 , CL or F and R_1 is H and when $R_1=R_2$, R_2 is OCH_3 or OCH_2O and R is H .

1/5



2, Combretastatin A-2

1a, R = OH, R₁ = OH

Combretastatin A-1

1b, R = OH, R₁ = H

Combretastatin A-4

1c, R = OSi(CH₃)₂C(CH₃)₃, R₁ = H1d, R = OPO₃Na₂, R₁ = H

Combretastatin A-4 prodrug

1e, R = R₁ = H3a, R = OSi(CH₃)₂C(CH₃)₃



3b, R = OH, Phenstatin

3c, R = OPO₃(C₆H₅CH₂)₂3d, R = OPO₃Na₂


Phenstatin prodrug

3e, R = OCOCH₃

3f, R = H


4a, R = H, R₁, R₂ = OCH₂O4b, R = R₂ = CH₃, R₁ = H4c, R = H, R₁ = R₂ = OCH₃4d, R = R₂ = OCH₃, R₁ = H4e, R = R₂ = Cl, R₁ = H4f, R = R₂ = F, R₁ = H

5



7a, R = OH

7b, R = H



8a, R = OH

8b, R = H

Figure 1.

09582952-090500